AMENDMENT AND RESPONSE TO OFFICE ACTION

APPENDIX II: CLEAN COPY OF CLAIMS AS PENDING

1. (amended) A compound of Formula I:

wherein

R¹, R², R³ and R⁴ are independently

H,

HO,

R¹³O-,

Halogen,

C1-C3-alkyl,

CF_{3.}

R14CO2-,

R14O2C-,

R¹⁴CO-,

R14CONH-,

R14NHCO-,

R¹⁴NHCO₂-,

R14OCONH-,

```
U.S.S.N.: 09/882,843
Filed: June 15, 2001
```

R14O2S-,

R14QS-,

R¹⁴S-, or

R15R16N-; or

R¹ and R², or R² and R³, or R³ and R⁴ taken together can be

-SCH₂S-,

-SCH₂O-,

-OCH₂S-,

-SCH2CH2S-,

-SCH₂CH₂O-, or

-OCH2CH2S-;

wherein one of R¹, R², R³ and R⁴ must be C1-C3-alkoxy or C1-C3-alkylthio group;

R⁵, R⁶, R⁷, and R⁸ are independently

H,

C1-C6-alkyl,

C3-C6-alkenyl,

C3-C6-cycloalkyl,

phenyl or substituted phenyl, wherein the phenyl is substituted with one or two

substituents, C1-C3-alkyl, halogen, R13O-, CF3-, R14O2S-, R14CO, R14CO2-,

R¹⁴O₂C-, R¹⁴CONH-, R¹⁴NHCO; or

R⁵ and R⁶ taken together can be C3-C6-cycloalkyl;

 R^7 and R^8 taken together can be C3-C6-cycloalkyl;

62

```
U.S.S.N.: 09/882,843
Filed: June 15, 2001
```

R⁹ is

R15R16NCO-,

R15R16NCS-,

R15R16N(CR17)-,

R¹⁷OCO-,

R¹⁵CO-,

R15R16NCH2CO-,

R¹⁴O₂C-(CH₂)_n-,

R¹⁵R¹⁶NCO-(CH₂)_n-,

 $NC-(CH_2)_{n-}$

H,

C1-C6-alkyl,

C3-C6-alkenyl, or

C3-C6-cycloalkyl; or

R8 and R9 taken together can be

 $-(CH_2)_mCH_2(R^{15})NCO-,$

-(CH₂)_mCH₂OCO-, or

-(CH₂)_mCH₂CH₂CO-;

 R^{10} and R^{11} are independently

H,

R15R16N-,

R¹⁵R¹⁶N(CR¹⁷)-,

63

```
U.S.S.N.: 09/882,843
Filed: June 15, 2001
```

R14HNCO-, or

R14CONH-;

R¹²is

H,

Halogen,

HO,

R¹³O-,

R15R16N-.

C1-C3-alkyl,

CF₃,

R14CO2-,

R14CO-, or

R14CONH-;

R¹³ is C1-C3-alkyl;

R¹⁴ is H or C1-C3-alkyl;

R¹⁵ and R¹⁶ are independently

Η,

C1-C10-alkyl,

C1-C6-perfluoroalkyl,

C3-C10-alkenyl, or

C3-C6-cycloalkyl; or

 R^{15} and R^{16} taken together can be C3-C6-cycloalkyl;

```
U.S.S.N.: 09/882,843
Filed: June 15, 2001
AMENDMENT AND RESPONSE TO OFFICE ACTION

R<sup>17</sup> is C1-C6-alkyl, C3-C6-alkenyl, or C3-C6-cycloalkyl;

n is 1 to 6;

m is 0 to 2;

and pharmaceutically acceptable salts thereof;

wherein R<sup>10</sup> and R<sup>11</sup> cannot be both H.

2. (amended) The compound of claim 1 of Formula I wherein
```

one of four substituents of R¹, R², R³ and R⁴ must be C1-C3-alkylthio group or C1-C3-alkoxy group, the other substituents are independently H, R¹³O-, R¹⁴S-, halogen[(F, Cl, Br)], or C1-C3-alkyl;

 R^2 and R^3 taken together can be $-SCH_2S$ -, $-SCH_2O$ -, or $-OCH_2S$ -; R^9 is

R¹⁵R¹⁶NCO-, R¹⁵R¹⁶NCS-, R¹⁵R¹⁶N(CR¹⁷)-, R¹⁷OCO-, or R¹⁵CO-

H;

R¹⁰ and R¹¹ are independently H, H₂N-, or CH₃CONH-; and pharmaceutically acceptable salts thereof.

3. (amended) A composition comprising the compound of claim 2 and a pharmaceutically acceptable carrier.

AMENDMENT AND RESPONSE TO OFFICE ACTION

- 4. (amended) The composition of claim 3 in a dosage form comprising a therapeutically effective amount of the compound for treating a disorder in a patient associated with excessive activation of the α-amino-3-hydroxy-5-methyl-4-isooxazoleproprionic acid (AMPA) subtype of the ionotropic excitatory amino acid (EAA) receptors.
- 5. (amended) The compound of claim 2 of Formula I selected from the group consisting of

1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-acetyl-7-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-methylcarbamoyl-7-methoxy-5H-2,3benzodiazepine, 1-(4-Aminophenyl)-3.5-dihydro-4-methyl-3-ethylcarbamoyl-7-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-propylcarbamoyl-7-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-butylcarbamoyl-7methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-3,5-dihydro-4-methyl-3-acetyl-7methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-3,5-dihydro-4-methyl-3methylcarbamoyl-7-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-3,5-dihydro-4-methyl-3-ethylcarbamoyl-7-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-3,5-dihydro-4-methyl-3-propylcarbamoyl-7-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-3,5-dihydro-4-methyl-3-butylcarbamoyl-7-methoxy-5H-2,3benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-acetyl-8-methoxy-5H-2,3benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-methylcarbamoyl-8-methoxy-5*H*-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-ethylcarbamoyl-8-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-propylcarbamoyl-8methoxy-5*H*-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-butylcarbamoyl-

AMENDMENT AND RESPONSE TO OFFICE ACTION

8-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-7-amino-3,5-dihydro-4-methyl-3-acetyl-8-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-7-amino-3,5-dihydro-4-methyl-3methylcarbamoyl-8-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-7-amino-3,5-dihydro-4-methyl-3-ethylcarbamoyl-8-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-7-amino-3,5-dihydro-4-methyl-3-propylcarbamoyl-8-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-7-amino-3,5-dihydro-4-methyl-3-butylcarbamoyl-8-methoxy-5H-2,3benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-acetyl-7-methylthio-5H-2,3benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-methylcarbamoyl-7-methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-ethylcarbamoyl-7methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3propylcarbamoyl-7-methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4methyl-3-butylcarbamoyl-7-methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-3,5-dihydro-4-methyl-3-acetyl-7-methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-8amino-3.5-dihydro-4-methyl-3-methylcarbamoyl-7-methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-3,5-dihydro-4-methyl-3-ethylcarbamoyl-7-methylthio-5H-2,3benzodiazepine, 1-(4-Aminophenyl)-8-amino-3,5-dihydro-4-methyl-3-propylcarbamoyl-7methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-3,5-dihydro-4-methyl-3butylcarbamoyl-7-methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4methyl-3-acetyl-8-methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4methyl-3-methylcarbamoyl-8-methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5dihydro-4-methyl-3-ethylcarbamoyl-8-methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-propylcarbamoyl-8-methylthio-5*H*-2,3-benzodiazepine, 1-(4-

67

AMENDMENT AND RESPONSE TO OFFICE ACTION

Aminophenyl)-3,5-dihydro-4-methyl-3-butylcarbamoyl-8-methylthio-5*H*-2,3-benzodiazepine, 1-(4-Aminophenyl)-7-amino-3,5-dihydro-4-methyl-3-acetyl-8-methylthio-5*H*-2,3-benzodiazepine, 1-(4-Aminophenyl)-7-amino-3,5-dihydro-4-methyl-3-methylcarbamoyl-8-methylthio-5*H*-2,3-benzodiazepine, 1-(4-Aminophenyl)-7-amino-3,5-dihydro-4-methyl-3-ethylcarbamoyl-8-methylthio-5*H*-2,3-benzodiazepine, 1-(4-Aminophenyl)-7-amino-3,5-dihydro-4-methyl-3-propylcarbamoyl-8-methylthio-5*H*-2,3-benzodiazepine, and 1-(4-Aminophenyl)-7-amino-3,5-dihydro-4-methyl-3-butylcarbamoyl-8-methylthio-5*H*-2,3-benzodiazepine.

- 6. (amended) A composition comprising the compound of claim 5 and a pharmaceutically acceptable carrier.
- 7. (amended) The composition of claim 6 in a dosage form comprising a therapeutically effective amount of the compound for treating a disorder in a patient associated with excessive activation of the α-amino-3-hydroxy-5-methyl-4-isooxazoleproprionic acid (AMPA) subtype of the ionotropic excitatory amino acid (EAA) receptors.
- 8. (amended) A composition comprising the compound of claim 1 and a pharmaceutically acceptable carrier.
- 9. (amended) The composition of claim 8 in a dosage form comprising a therapeutically effective amount of the compound for treating a disorder in a patient associated with excessive activation of the α-amino-3-hydroxy-5-methyl-4-isooxazoleproprionic acid (AMPA) subtype of the ionotropic excitatory amino acid (EAA) receptors.
- 10. (amended) A method for treating a patient having a disorder associated with excessive activation of the α-amino-3-hydroxy-5-methyl-4-isooxazoleproprionic acid (AMPA) subtype of the ionotropic excitatory amino acid (EAA) receptors, the method comprising

68

AMENDMENT AND RESPONSE TO OFFICE ACTION

administering to the patient, in an effective amount to alleviate the symptoms of the disorder, a

compound of Formula I:

wherein

R¹, R², R³ and R⁴ are independently

H,

HO,

· R13O-.

halogen,

C1-C3-alkyl,

CF_{3,}

R¹⁴CO_{2*},

 $R^{14}O_2C_{-}$

R14CO-,

R¹⁴CONH-,

R14NHCO-,

R¹⁴NHCO₂-,

R14OCONH-,

```
U.S.S.N.: 09/882,843
Filed: June 15, 2001
```

R14O2S-,

R¹⁴OS-.

R¹⁴S-. or

R¹⁵R¹⁶N-: or

R¹ and R², or R² and R³, or R³ and R⁴ taken together can be

-SCH₂S-,

-SCH₂O-,

-OCH₂S-,

-SCH₂CH₂S-,

-SCH₂CH₂O-, or

-OCH2CH2S-;

wherein one of R¹, R², R³ and R⁴ must be C1-C3-alkoxy or C1-C3-alkylthio group;

R⁵, R⁶, R⁷, and R⁸ are independently

H,

C1-C6-alkyl,

C3-C6-alkenyl,

C3-C6-cycloalkyl,

phenyl or substituted phenyl, wherein the phenyl is substituted with one or two

substituents, C1-C3-alkyl, halogen, R13O-, CF3-, R14O2S-, R14CO, R14CO2-,

R¹⁴O₂C-, R¹⁴CONH-, R¹⁴NHCO; or

 R^5 and R^6 taken together can be C3-C6-cycloalkyl;

R⁷ and R⁸ taken together can be C3-C6-cycloalkyl;

70

```
U.S.S.N.: 09/882,843
Filed: June 15, 2001
```

R⁹ is

R15R16NCO-,

R15R16NCS-.

. R15R16N(CR17)-,

R¹⁷OCO-,

R¹⁵CO-,

R¹⁵R¹⁶NCH₂CO-,

 $R^{14}O_2C-(CH_2)_{n}$ -,

R¹⁵R¹⁶NCO-(CH₂)_n-,

 $NC-(CH_2)_{n-}$

H,

C1-C6-alkyl,

C3-C6-alkenyl, or

C3-C6-cycloalkyl; or

R8 and R9 taken together can be

 $-(CH_2)_mCH_2(R^{15})NCO-,$

-(CH₂)_mCH₂OCO-, or

-(CH₂)_mCH₂CH₂CO-;

R¹⁰ and R¹¹ are independently

H,

 $R^{15}R^{16}N$ -,

 $R^{15}R^{16}N(CR^{17})$ -,

71

```
U.S.S.N.: 09/882,843
Filed: June 15,2001
```

R¹⁴HNCO-, or

R14CONH-;

R¹² is

H,

Halogen,

HO,

R13O-.

 $R^{15}R^{16}N_{-}$

C1-C3-alkyl,

CF₃,

R14CO2-,

R¹⁴CO-, or

R14CONH-:

R¹³ is C1-C3-alkyl;

R¹⁴ is H or C1-C3-alkyl;

R¹⁵ and R¹⁶ are independently

H,

C1-C10-alkyl,

C1-C6-perfluoroalkyl,

C3-C10-alkenyl, or

C3-C6-cycloalkyl; or

R¹⁵ and R¹⁶ taken together can be C3-C6-cycloalkyl;

72

```
U.S.S.N.: 09/882,843
Filed: June 15, 2001
```

R¹⁷ is C1-C6-alkyl, C3-C6-alkenyl, or C3-C6-cycloalkyl;

n is 1 to 6;

m is 0 to 2:

and pharmaceutically acceptable salts thereof;

wherein R¹⁰ and R¹¹ cannot be both H.

in combination with a pharmaceutically acceptable carrier.

11. (amended) The method of claim 10 wherein, in the compound of Formula I, one of four substituents of R¹, R², R³ and R⁴ must be C1-C3-alkylthio group or C1-C3-alkoxy group, the other substituents are independently H, R¹³O-, R¹⁴S-, halogen, or C1-C3-alkyl; R² and R³ taken together can be -SCH₂S-, -SCH₂O-, or -OCH₂S-; R⁹ is

R15R16NCO-,

R15R16NCS-,

R15R16N(CR17)-.

R¹⁷OCO-, or

R15CO-,

H;

R¹⁰ and R¹¹ are independently H, H₂N-, or CH₃CONH-; and pharmaceutically acceptable salts thereof.


12. The method of claim 11 wherein the disorder is selected from the group consisting of neurological, neuropsychological, neuropsychological, neuropsychopharmacological and functional disorders.

73

AMENDMENT AND RESPONSE TO OFFICE ACTION

13. (amended) The method of claim 11 wherein the compound of Formula I is selected ... from the group consisting of

1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-acetyl-7-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-methylcarbamoyl-7-methoxy-5*H*-2,3benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-ethylcarbamoyl-7-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-propylcarbamoyl-7-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-butylcarbamoyl-7methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-3,5-dihydro-4-methyl-3-acetyl-7methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-3,5-dihydro-4-methyl-3methylcarbamoyl-7-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-3,5-dihydro-4-methyl-3-ethylcarbamoyl-7-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-3,5-dihydro-4-methyl-3-propylcarbamoyl-7-methoxy-5H-2.3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-3,5-dihydro-4-methyl-3-butylcarbamoyl-7-methoxy-5H-2,3benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-acetyl-8-methoxy-5H-2,3benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-methylcarbamoyl-8-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-ethylcarbamoyl-8-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-propylcarbamoyl-8methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-butylcarbamoyl-8-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-7-amino-3,5-dihydro-4-methyl-3-acetyl-8-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-7-amino-3,5-dihydro-4-methyl-3methylcarbamoyl-8-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-7-amino-3,5-dihydro-4-methyl-3-ethylcarbamoyl-8-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-7-amino-



AMENDMENT AND RESPONSE TO OFFICE ACTION

3,5-dihydro-4-methyl-3-propylcarbamoyl-8-methoxy-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-7-amino-3,5-dihydro-4-methyl-3-butylcarbamoyl-8-methoxy-5H-2,3benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-acetyl-7-methylthio-5H-2,3benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-methylcarbamoyl-7-methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-ethylcarbamoyl-7methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3propylcarbamoyl-7-methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4methyl-3-butylcarbamoyl-7-methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-3,5-dihydro-4-methyl-3-acetyl-7-methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-8amino-3,5-dihydro-4-methyl-3-methylcarbamoyl-7-methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-3,5-dihydro-4-methyl-3-ethylcarbamoyl-7-methylthio-5H-2,3benzodiazepine, 1-(4-Aminophenyl)-8-amino-3,5-dihydro-4-methyl-3-propylcarbamoyl-7methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-3,5-dihydro-4-methyl-3butylcarbamoyl-7-methylthio-5*H*-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4methyl-3-acetyl-8-methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4methyl-3-methylcarbamoyl-8-methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5dihydro-4-methyl-3-ethylcarbamoyl-8-methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-propylcarbamoyl-8-methylthio-5*H*-2,3-benzodiazepine, 1-(4-Aminophenyl)-3,5-dihydro-4-methyl-3-butylcarbamoyl-8-methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-7-amino-3,5-dihydro-4-methyl-3-acetyl-8-methylthio-5H-2,3-benzodiazepine, 1-(4-Aminophenyl)-7-amino-3,5-dihydro-4-methyl-3-methylcarbamoyl-8-methylthio-5H-2.3benzodiazepine, 1-(4-Aminophenyl)-7-amino-3,5-dihydro-4-methyl-3-ethylcarbamoyl-8-

AMENDMENT AND RESPONSE TO OFFICE ACTION

methylthio-5*H*-2,3-benzodiazepine, 1-(4-Aminophenyl)-7-amino-3,5-dihydro-4-methyl-3-propylcarbamoyl-8-methylthio-5*H*-2,3-benzodiazepine, and 1-(4-Aminophenyl)-7-amino-3,5-dihydro-4-methyl-3-butylcarbamoyl-8-methylthio-5*H*-2,3-benzodiazepine.

- 14. The method of claim 13 wherein the disorder is selected from the group consisting of neurological, neuropsychological, neuropsychiatric, neurodegenerative, neuropsychopharmacological and functional disorders.
- 15. The method of claim 10 wherein the disorder is selected from the group consisting of neurological, neuropsychological, neuropsychological, neuropsychopharmacological and functional disorders.
 - 16. (amended) A compound of Formula II:

wherein

R¹ and R⁴ are independently

H,

HO,

R13O-.

Halogen,

C1-C3-alkyl,

76

9YM (14 77588/3

AMENDMENT AND RESPONSE TO OFFICE ACTION

CF_{3.}

R14CO2-,

 $R^{14}O_2C_{-}$

R¹⁴CO-.

R¹⁴CONH-.

R¹⁴NHCO-,

R¹⁴NHCO₂-,

R14OCONH-.

R¹⁴O₂S-,

R14OS-,

R14S-, or

R¹⁵R¹⁶N-; or

R² is one of H, HO, R¹³O-, halogen, C1-C3-alkyl, CF₃, R¹⁴CO₂-, R¹⁴O₂C-, R¹⁴CO-, R¹⁴CONH-, R¹⁴NHCO-, R¹⁴NHCO₂-, R¹⁴OCONH-, R¹⁴O₂S-, R¹⁴OS-, R¹⁴S- and R¹⁵R¹⁶N- when R³ is one of HO, halogen, C1-C3-alkyl, CF₃, R¹⁴CO₂-, R¹⁴O₂C-, R¹⁴CO-, R¹⁴CONH-, R¹⁴NHCO-, R¹⁴NHCO₂-, R¹⁴OCONH-, R¹⁴O₂S-, R¹⁴OS-, R¹⁴S-, and R¹⁵R¹⁶N-; or

R² is one of H, HO, halogen, C1-C3-alkyl, CF₃, R¹⁴CO₂-, R¹⁴O₂C-, R¹⁴CO-, R¹⁴CO-, R¹⁴CONH-, R¹⁴NHCO-, R¹⁴NHCO₂-, R¹⁴OCONH-, R¹⁴O₂S-, R¹⁴OS-, R¹⁴S- and R¹⁵R¹⁶N- when R³ is one of H, HO, R¹³O-, halogen, C1-C3-alkyl, CF₃, R¹⁴CO₂-, R¹⁴O₂C-, R¹⁴CO-, R¹⁴CONH-, R¹⁴NHCO-, R¹⁴NHCO₂-, R¹⁴OCONH-, R¹⁴O₂S-, R¹⁴OS-, R¹⁴S-, and R¹⁵R¹⁶N-; or

R¹ and R², or R² and R³, or R³ and R⁴ taken together can be -SCH₂S-,

77

AMENDMENT AND RESPONSE TO OFFICE ACTION

- -SCH₂O-,
- -OCH₂S-,
- -SCH₂CH₂S-,
- -SCH₂CH₂O-, or
- -OCH₂CH₂S-; or

one of four substituents of R¹, R², R³ and R⁴ must be C1-C3-alkoxy or C1-C3-alkylthio group;

R⁵, R⁶, and R⁷ are independently

H,

C1-C6-alkyl,

C3-C6-alkenyl,

C3-C6-cycloalkyl, or

phenyl or substituted phenyl, wherein the phenyl is substituted with one or two

substituents, C1-C3-alkyl, halogen, R¹³O-, CF₃-, R¹⁴O₂S-, R¹⁴OS-, R¹⁴CO, R¹⁴CO₂-,

R¹⁴O₂C-, R¹⁴CONH-, R¹⁴NHCO; or

R⁵ and R⁶ taken together can be C3-C6-cycloalkyl;

R¹³ is C1-C3-alkyl;

R¹⁴ is H or C1-C3-alkyl;

 R^{15} and R^{16} are independently

H,

C1-C10-alkyl,

C1-C6-perfluoroalkyl,

78

AMENDMENT AND RESPONSE TO OFFICE ACTION

C3-C10-alkenyl, or

C3-C6-cycloalkyl; or

R¹⁵ and R¹⁶ taken together can be C3-C6-cycloalkyl;

R¹⁸ and R¹⁹ are independently

H,

Halogen,

C1-C3-alkyl,

R14O-,

CF₃-, or

 $R^{14}CO_2$ -;

R²⁰ and R²¹ are independently

H,

R15R16N-,

R¹⁵HNC(NH)-, or

R14CONH-;

and pharmaceutically acceptable salts thereof;

wherein R²⁰ and R²¹ cannot both be H.

17. (amended) The compound of claim 16 of Formula II wherein one of four substituents of R¹, R², R³ and R⁴ must be C1-C3-alkylthio or C1-C3-alkoxy group, the other substituents are independently H, R¹³O-, R¹³S-, halogen, or C1-C3-alkyl; R² and R³ taken together can be -SCH₂S-, -SCH₂O-, or -OCH₂S-;

79

AMENDMENT AND RESPONSE TO OFFICE ACTION

R²⁰ and R²¹ are independently H, H₂N-, or CH₃CONH-; and pharmaceutically acceptable salts thereof.

- 18. (amended) A composition comprising the compound of claim 17 and a pharmaceutically acceptable carrier.
- 19. (amended) The composition of claim 18 in a dosage form comprising a therapeutically effective amount of the compound for treating a disorder in a patient associated with excessive activation of the α-amino-3-hydroxy-5-methyl-4-isooxazoleproprionic acid (AMPA) subtype of the ionotropic excitatory amino acid (EAA) receptors.
- 20. (amended) The compound of claim 17 of Formula II selected from the group consisting of

1-(4-Aminophenyl)-4-methyl-7-methoxy-5*H*-2,3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-4-methyl-7-methoxy-5*H*-2,3-benzodiazepine, 1-(4-Aminophenyl)-7-amino-4-methyl-8-methoxy-5*H*-2,3-benzodiazepine, 1-(4-Aminophenyl)-4-methyl-7-methylthio-5*H*-2,3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-4-methyl-7-methylthio-5*H*-2,3-benzodiazepine, 1-(4-Aminophenyl)-4-methyl-8-methylthio-5*H*-2,3-benzodiazepine, and 1-(4-Aminophenyl)-7-amino-4-methyl-8-methylthio-5*H*-2,3-benzodiazepine.

- 21. (amended) A composition comprising the compound of claim 20 and a pharmaceutically acceptable carrier.
- 22. (amended) The composition of claim 21 in a dosage form comprising a therapeutically effective amount of the compound for treating a disorder in a patient associated with excessive activation of the α-amino-3-hydroxy-5-methyl-4-isooxazoleproprionic acid (AMPA) subtype of the ionotropic excitatory amino acid (EAA) receptors.

80

AMENDMENT AND RESPONSE TO OFFICE ACTION

- 23. (amended) A composition comprising the compound of claim 16 and a pharmaceutically acceptable carrier.
- 24. (amended) The composition of claim 23 in a dosage form comprising a therapeutically effective amount of the compound for treating a disorder in a patient associated with excessive activation of the α-amino-3-hydroxy-5-methyl-4-isooxazoleproprionic acid (AMPA) subtype of the ionotropic excitatory amino acid (EAA) receptors.
- 25. (amended) A method for treating a patient having a disorder associated with excessive activation of the α-amino-3-hydroxy-5-methyl-4-isooxazoleproprionic acid (AMPA) subtype of the ionotropic excitatory amino acid (EAA) receptors, the method comprising administering to the patient, in an effective amount to alleviate the symptoms of the disorder, a compound of Formula II:

wherein

 R^1 and R^4 are independently

H.

HO,

R13O-,

Halogen,

81

AMENDMENT AND RESPONSE TO OFFICE ACTION

C1-C3-alkyl,

CF₃

 $R^{14}CO_{27}$

 $R^{14}O_2C_{-}$

R¹⁴CO-.

R¹⁴CONH-,

R¹⁴NHCO-.

R14NHCO2-,

R¹⁴OCONH-,

 $R^{14}O_2S_{-}$

R¹⁴OS-,

R14S-, or

R15R16N-; or

 R^2 is one of H, HO, R^{13} O-, halogen, C1-C3-alkyl, CF₃, R^{14} CO₂-, R^{14} O₂C-, R^{14} CO-, R^{14} CONH-, R^{14} NHCO-, R^{14} NHCO₂-, R^{14} OCONH-, R^{14} OCONH-, R^{14} OS-, R^{14} S- and R^{15} R¹⁶N- when R^3 is one of HO, halogen, C1-C3-alkyl, CF₃, R^{14} CO₂-, R^{14} O₂C-, R^{14} CO-, R^{14} CONH-, R^{14} NHCO-, R^{14} NHCO-, R^{14} OCONH-, R^{14} O2S-, R^{14} OS-, R^{14} S-, and R^{15} R¹⁶N-; or

R² is one of H, HO, halogen, C1-C3-alkyl, CF₃, R¹⁴CO₂-, R¹⁴O₂C-, R¹⁴CO-, R¹⁴CONH-, R¹⁴NHCO-, R¹⁴NHCO₂-, R¹⁴OCONH-, R¹⁴O₂S-, R¹⁴OS-, R¹⁴S- and R¹⁵R¹⁶N- when R³ is one of H, HO, R¹³O-, halogen, C1-C3-alkyl, CF₃, R¹⁴CO₂-, R¹⁴O₂C-, R¹⁴CO-, R¹⁴CONH-, R¹⁴NHCO-, R¹⁴NHCO₂-, R¹⁴OCONH-, R¹⁴O₂S-, R¹⁴OS-, R¹⁴S-, and R¹⁵R¹⁶N-; or

R¹ and R², or R² and R³, or R³ and R⁴ taken together can be

82

AMENDMENT AND RESPONSE TO OFFICE ACTION

-SCH₂S-,

-SCH₂O-,

-OCH₂S-,

-SCH₂CH₂S-,

-SCH₂CH₂O-, or

-OCH2CH2S-; or

one of four substituents of R¹, R², R³ and R⁴ must be C1-C3-alkoxy or C1-C3-alkylthio group;

R⁵, R⁶, and R⁷ are independently

H,

C1-C6-alkyl,

C3-C6-alkenyl,

C3-C6-cycloalkyl, or

phenyl or substituted phenyl, wherein the phenyl is substituted with one or two

substituents, C1-C3-alkyl, halogen, R13O-, CF3-, R14O2S-, R14CO, R14CO2-,

R¹⁴O₂C-, R¹⁴CONH-, R¹⁴NHCO; or

R⁵ and R⁶ taken together can be C3-C6-cycloalkyl;

R¹³ is C1-C3-alkyl;

R¹⁴ is H or C1-C3-alkyl;

R¹⁵ and R¹⁶ are independently

H.

C1-C10-alkyl,

83

AMENDMENT AND RESPONSE TO OFFICE ACTION

C1-C6-perfluoroalkyl,

C3-C10-alkenyl, or

C3-C6-cycloalkyl; or

 R^{15} and R^{16} taken together can be C3-C6-cycloalkyl;

R¹⁸ and R¹⁹ are independently

H,

Halogen,

C1-C3-alkyl,

R14O-.

CF₃-, or

R14CO2-;

R²⁰ and R²¹ are independently

H,

R15R16N-,

R15HNC(NH)-, or

R¹⁴CONH-:

and pharmaceutically acceptable salts thereof;

wherein R²⁰ and R²¹ cannot both be H,

in combination with a pharmaceutically acceptable carrier.

26. (amended) The method of claim 25 wherein, in the compound of Formula II, one of four substituents of R¹, R², R³ and R⁴ must be C1-C3-alkylthio or C1-C3-alkoxy group, the other substituents are independently H, R¹³O-, R¹³S-, halogen, or C1-C3-alkyl;

84

AMENDMENT AND RESPONSE TO OFFICE ACTION

R² and R³ taken together can be -SCH₂S-, -SCH₂O-, or -OCH₂S-;

R²⁰ and R²¹ are independently H, H₂N-, or CH₃CONH-; and pharmaceutically acceptable salts thereof.

- 27. The method of claim 26 wherein the disorder is selected from the group consisting of neurological, neuropsychological, neuropsychiatric, neurodegenerative, neuropsychopharmacological and functional disorders.
- 28. (amended) The method of claim 26 wherein the compound of Formula II is selected from the group consisting of

1-(4-Aminophenyl)-4-methyl-7-methoxy-5*H*-2,3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-4-methyl-7-methoxy-5*H*-2,3-benzodiazepine, 1-(4-Aminophenyl)-4-methyl-7-amino-4-methyl-8-methoxy-5*H*-2,3-benzodiazepine, 1-(4-Aminophenyl)-4-methyl-7-methylthio-5*H*-2,3-benzodiazepine, 1-(4-Aminophenyl)-8-amino-4-methyl-7-methylthio-5*H*-2,3-benzodiazepine, 1-(4-Aminophenyl)-7-amino-4-methyl-8-methylthio-5*H*-2,3-benzodiazepine, and 1-(4-Aminophenyl)-7-amino-4-methyl-8-methylthio-5*H*-2,3-benzodiazepine,

- 29. The method of claim 28 wherein the disorder is selected from the group consisting of neurological, neuropsychological, neuropsychological, neuropsychological, neuropsychological and functional disorders.
- 30. The method of claim 25 wherein the disorder is selected from the group consisting of neurological, neuropsychological, neuropsychiatric, neurodegenerative, neuropsychopharmacological and functional disorders.

ATL1 #568404 v1